

Thermodynamical Control by Frequent Quantum Measurements*

Noam Erez, Goren Gordon, and Gershon Kurizki†

Department of Chemical Physics, Weizmann Institute of Science, Rehovot 76100, Israel.

Mathias Nest

Theoretische Chemie, Universitaet Potsdam, Potsdam 14476, Germany.

Heat flow between a large “bath” and a smaller system brings them progressively closer to thermal equilibrium while increasing their entropy[1]. Deviations from this trend are fluctuations involving a small fraction of a statistical ensemble of systems interacting with the bath: in this respect, quantum and classical thermodynamics are in agreement[1, 2, 3, 4, 5]. Can there be drastic differences between them? Here we address a distinctly quantum mechanical setting that displays such differences: disturbances of thermal equilibrium between two-level systems (TLS) and a bath[6] by frequent and brief quantum (non-demolishing)[7, 8, 9, 10] measurements of the TLS energy-states. If the measurements are frequent enough to induce either the Zeno or the anti-Zeno regime, namely, the slowdown or speedup of the TLS relaxation[11, 12, 13, 14, 15], then the resulting entropy and temperature of both the system and the bath are found to be completely unrelated to what is expected by standard thermodynamical rules that hold for memoryless baths[2, 5]. The practical advantage of these anomalies is the possibility of *very fast control* of heat and entropy, allowing cooling and state-purification of quantum systems much sooner than their thermal equilibration time.

To understand the origins of the predicted anomalies, consider a thermal bath in equilibrium with an ensemble of quantum systems. The energy of the quantum systems is briefly measured. How will these systems be affected? Classically, their equilibrium state may remain intact, since measurements can be chosen to be non-intrusive, i.e., involve no energy exchange, but merely provide “snapshots” of the system. Likewise, quantum mechanically, nearly-ideal (projective) measurements involve no energy cost when performed by macroscopic detectors on isolated systems[8]. Yet finite-time coupling, followed by abrupt decoupling, of two quantum ensembles, which may be viewed as the detection of one ensemble by the other, may cause an increase of their mean total energy[16]. Here we address a different scenario pertaining to thermodynamics: a detector briefly (nearly-impulsively) measures only two-level systems (TLS) that are initially at thermal equilibrium with a much larger bath. We then ask: how will the temperature and entropy of these systems evolve both during and after the measurement via an interplay between the detector, the system and the bath?

Such nearly-impulsive quantum measurements in the basis of the system (energy) eigenstates, chosen to be the “pointer basis”[7] of the detector, must transfer energy, via the detector-system coupling, so as to momentarily interrupt (override) the system-bath interaction. This energy transfer, resulting in a change in system-bath entanglement, triggers the distinctly quantum dynamics of both the system and the bath, which subsequently re-

distributes their mean energy and entropy in anomalous, unfamiliar, ways.

It is possible to use the information gained by such measurements to sort out system subensembles according to their measured energy, in order to extract work or entropy change, in the spirit of Maxwell’s demon[17]. Here, however, we let the entire TLS ensemble evolve regardless of the measured result, i.e., we trace out the detector states. Our main concern is with the question: if we probe this evolution by a subsequent measurement, how will the outcome depend on the time-separation of the two? The answer should elucidate the virtually unknown short-time evolution of quantum systems coupled to a bath.

Specifically, we analyze the scenario described above for a TLS with energy separation $\hbar\omega_a$, that is weakly coupled to a thermal bath of harmonic oscillators, characterized by a correlation (memory) time of the bath response $t_c \gg 1/\omega_a$, which typically marks the onset of equilibrium. After equilibrium has been reached, we perform $k = 1, \dots, K$ quantum non-demolition (QND) measurements[7, 8, 9, 10] of the TLS energy states at times separated by $\Delta t_k = t_{k+1} - t_k$. Each measurement has a brief duration $\tau_k \ll 1/\omega_a$. Our aim is to explore the evolution as a function of time-separations between consecutive measurements in the non-Markov domain, $\Delta t_k \lesssim 1/\omega_a \ll t_c$. Such measurements do not resolve the energies of the TLS states, due to the time-energy uncertainty. Yet they can discriminate between states of different *symmetry*, e.g., different angular momenta. In a TLS, the evolution of the mean energy or state populations can be identified with the effective (spin) temperature change[18], however rapidly it occurs. In this uncharted domain, we show that consecutive brief measurements entail several anomalies: (i) The quantum-mechanical non-commutativity of the system-

*This is the presubmission version of *Nature* **452**, 724. The published version is available on the *Nature* website: <http://dx.doi.org/10.1038/nature06873>.

†Electronic address: gershon.kurizki@weizmann.ac.il

detector and system-bath interactions causes the heat-up of the system at the expense of the detector-system coupling, but not at the expense of the coupling to the bath, only at very short Δt_k compatible with the quantum Zeno effect (QZE)[11]. (ii) A *transition from heating to cooling* of the TLS ensemble may occur as we vary the interval between consecutive measurements from $\Delta t_k \ll 1/\omega_a$ to $\Delta t_k \sim 1/\omega_a \ll t_c$. This marks the transition from Δt_k compatible with the QZE to those compatible with the anti-Zeno effect (AZE)[12, 13, 14, 15]. Remarkably, the cooling may occur even if the bath is initially hotter. (iii) Correspondingly, *oscillations of the entropy* relative to that of the equilibrium state take place, contrary to the Markovian notion of the second law of thermodynamics[2, 5].

This scenario is governed by the following total Hamiltonian of the system that interacts with the bath and is intermittently perturbed by the coupling of the system to the detector (measuring apparatus):

$$H(t) = H_{tot} + H_{SD}(t), \quad H_{tot} = H_S + H_B + H_{SB}. \quad (1)$$

Here H_S is the Hamiltonian of the TLS, with ground and excited states $|g\rangle, |e\rangle$, respectively; H_B is that of the thermal-bath composed of harmonic oscillators with energies $\hbar\omega_\lambda$; $H_{SB} = \mathcal{S}\mathcal{B}$ is the system-bath interaction Hamiltonian[18] (the spin-boson interaction): a product of the system-dipole (or spin-flip) operator $\mathcal{S}$ and the operator $\mathcal{B}$ describing the bath excitations and deexcitations; and $H_{SD}(t)$ is the time-dependent measurement Hamiltonian that couples the system to a detector comprised of energy-degenerate ancillae (for details see Supplement A). In the coupling Hamiltonians (H_{SB}, H_{SD}) we *do not* invoke the rotating-wave approximation (RWA)[18], namely, we do not impose energy conservation between the system and the bath or the detector, on the time scales considered[15].

The near-equilibrium state, ρ_{tot} , prior to a measurement has several pertinent characteristics (Supplement B): (a) It displays system-bath entanglement with off-diagonal matrix elements $\langle e|\rho_{tot}|g\rangle \neq 0$. (b) The system is described by a *diagonal* reduced density matrix, $\rho_S = \text{Tr}_B \rho_{tot}$, in the H_S eigenbasis. (c) The mean interaction energy $\langle H_{SB} \rangle$ is *negative*, assuming ρ_{tot} weakly deviates from the ground state of H_{tot} : $\langle H_{SB} \rangle = \langle H_{tot} \rangle - \langle H_S + H_B \rangle < 0$. This comes about since the correction to the ground-state energy of H_{tot} due to a weakly-perturbing interaction H_{SB} is negative (to the leading second order).

We next consider the disturbance of this equilibrium state by a nearly-impulsive (projective) quantum measurement ($\tau \rightarrow 0$) of S , in the $|g\rangle, |e\rangle$ basis. The measurement correlates the TLS energy eigenstates with mutually orthogonal states of an ancillary detector and the latter is then averaged (traced) over. This measurement has distinctly quantum-mechanical consequences (Supplement A): it interrupts the system-bath interaction[13], using the energy supplied by $H_{SD}(0 < t < \tau)$ (the system-detector coupling) without changing $\langle H_D \rangle$. It

thus eliminates the mean system-bath interaction energy, whose pre-measurement value was negative, as argued above:

$$\langle H_{SB}(0) \rangle < 0 \mapsto \langle H_{SB}(\tau) \rangle = 0, \quad \langle H_{SD}(t) \rangle = -\langle H_{SB}(t) \rangle. \quad (2)$$

We describe the detection process as a CNOT operation that retains the energy degeneracy of the detector $\langle H_D \rangle = 0$, although its Von-Neumann entropy increases (Supplement A).

After the measurement (as $H_{SD}(t \geq \tau) = 0$), time-energy uncertainty at $\Delta t \lesssim 1/\omega_a$ results in the breakdown of the RWA, i.e., $\langle H_S + H_B \rangle$ is not conserved as Δt grows. Only $\langle H_{tot} \rangle$ is conserved, by unitarity, until the next measurement. Hence, the post-measurement *decrease* of $\langle H_{SB} \rangle$ with Δt , signifying the restoration of equilibrium: $\langle H_{SB}(\tau) \rangle = 0 \rightarrow \langle H_{SB}(\tau + \Delta t) \rangle < 0$, is at the expense of an *increase* of $\langle H_S + H_B \rangle = \langle H_{tot} \rangle - \langle H_{SB} \rangle$, i.e., *heating* of the system and the bath, combined:

$$\left. \frac{d}{dt} [\langle H_S \rangle + \langle H_B \rangle] \right|_{\tau+\Delta t} > 0, \quad \left. \frac{d}{dt} \langle H_{SB} \rangle \right|_{\tau+\Delta t} < 0. \quad (3)$$

The post-measurement evolution of the system alone, described by $\rho_S = \text{Tr}_B \rho_{tot}$, is not at all obvious. Its Taylor expansion holds at short evolution times, $\Delta t \ll 1/\omega_a$,

$$\rho_S(\tau + \Delta t) \simeq \rho_S(\tau) + \Delta t \dot{\rho}_S(\tau) + \frac{\Delta t^2}{2} \ddot{\rho}_S(\tau) + \dots \quad (4)$$

The 0th order term is *unchanged* by the measurement, $\rho_S(\tau) = \rho_S(t \leq 0)$. The first derivative *vanishes* at $t = \tau$ ($\Delta t = 0$) due to the definite parity of the bath density-operator correlated to $|g\rangle$ or $|e\rangle$ (Supplement B). This initial post-measurement vanishing, $\dot{\rho}_S(\tau) = 0$, is the QZE condition[11, 13, 14, 15]. The time evolution of ρ_S is then governed by its second time derivative $\ddot{\rho}_S(\tau)$, which can be shown (Supplement B) to have the same sign as $\sigma_z = |e\rangle\langle e| - |g\rangle\langle g|$, the population difference operator of the TLS. Hence, the second derivative in (4) is *positive* shortly after the measurement, consistently with Eq. (3), if there is no initial population inversion of the system, i.e., for non-negative temperature.

The evolution of ρ_S at longer times (in the regime of weak system-bath coupling) may be approximately described (as verified by our *exact* numerical simulations[19], Supplement C) by the second-order non-Markovian master equation (ME) (Fig. 1 – main panel). The ME for ρ_S , on account of its diagonality, can be cast into the following population rate equations[15], dropping the subscript S in what follows and setting the measurement time to be $t = 0$:

$$\dot{\rho}_{ee}(t) = -\dot{\rho}_{gg}(t) = R_g(t)\rho_{gg} - R_e(t)\rho_{ee}, \quad (5)$$

$$R_{e(g)}(t) = 2\pi t \int_{-\infty}^{\infty} d\omega G_T(\omega) \text{sinc}[(\omega \mp \omega_a)t], \quad (6)$$

$$G_T(\omega) = (n_T(\omega) + 1)G_0(\omega) + n_T(-\omega)G_0(-\omega). \quad (7)$$

Here $\text{sinc}(x) = \frac{\sin(x)}{x}$, $G_T(\omega)$ is the temperature-dependent coupling spectrum of the bath, $G_0(\omega)$ is the zero-temperature coupling spectrum with peak coupling strength at ω_0 and spectral width $\sim 1/t_c$ and $n_T(\omega) = \frac{1}{e^{\beta\omega} - 1}$ is the inverse-temperature- (β -) dependent population of bath mode ω .

The entire dynamics is determined by $R_{e(g)}(t)$, the relaxation rates of the excited (ground) states. Their non-Markov time-dependence yields three distinct regimes:

(i) At short times $t \ll 1/\omega_a \ll t_c$ the *sinc* function in (6) is much broader than G_T . The relaxation rates R_e and R_g are then equal at any temperature, indicating the complete breakdown of the RWA discussed above: $|g\rangle \rightarrow |e\rangle$ and $|e\rangle \rightarrow |g\rangle$ transitions do not require quantum absorption or emission by the bath, respectively. The rates $R_{e(g)}$ then become *linear* in time, manifesting the QZE[13, 14, 15]:

$$R_{e(g)}(t \ll t_c) \approx 2\dot{R}_0 t, \quad (8)$$

$$\dot{R}_0 \equiv \int_{-\infty}^{\infty} d\omega G_T(\omega) = \langle B^2 \rangle. \quad (9)$$

This short-time regime implies the *universal Zeno heating rate*:

$$\frac{d}{dt}(\rho_{ee} - \rho_{gg}) \approx 4\dot{R}_0 t(\rho_{gg} - \rho_{ee}). \quad (10)$$

(ii) At intermediate non-Markovian times, $t \sim 1/\omega_a$, when the *sinc* function and G_T in (6) have comparable widths, the relaxation rates $R_{e(g)}(t)$ exhibit several unusual phenomena that stem from time-energy uncertainty. The change in the overlap of the *sinc* and G_T functions with time results in damped aperiodic oscillations of $R_e(t)$ and $R_g(t)$, near the frequencies $\omega_0 - \omega_a$ and $\omega_0 + \omega_a$, respectively. This oscillatory time dependence that conforms neither to QZE nor to the converse AZE of relaxation speedup[12, 13, 14], will henceforth be dubbed the *oscillatory Zeno effect* (OZE). Due to the negativity of the *sinc* function between its consecutive maxima, we can have a *negative relaxation rate*, which is completely forbidden by the RWA. Since $\text{sinc}[(\omega + \omega_a)t]$ is much further shifted from the peak of $G_T(\omega)$ than $\text{sinc}[(\omega - \omega_a)t]$, $R_g(t)$ is more likely to be negative than $R_e(t)$ (Fig. 1(a), Fig. 2(a)). Hence, $\rho_{gg}(t)$ may grow at the expense of $\rho_{ee}(t)$ more than allowed by the thermal-equilibrium detailed balance. This may cause *transient cooling*, as detailed below.

(iii) At long times $t \gg t_c$, the relaxation rates attain their Golden-Rule (Markov) values[15]

$$R_{e(g)}(t \gg t_c) \simeq 2\pi G_T(\pm\omega_a). \quad (11)$$

The populations then approach those of an equilibrium Gibbs state whose temperature is equal to that of the thermal bath (Fig. 1 – main panel).

We now turn to entropy dynamics. One may always define the entropy of ρ_S relative to its equilibrium state

ρ_0 (“entropy distance”) and the negative of its rate of change, as[3, 5]:

$$\mathcal{S}(\rho_S(t)||\rho_0) \equiv \text{Tr}\{\rho_S(t) \ln \rho_S(t)\} - \text{Tr}\{\rho_S(t) \ln \rho_0\} \quad (12a)$$

$$\sigma(t) \equiv -\frac{d}{dt}\mathcal{S}(\rho_S(t)||\rho_0). \quad (12b)$$

Only in the Markovian realm, σ thus defined is identified as the “entropy production rate”[2, 3, 5], where $\sigma(t) \geq 0$ is a statement of the second law of thermodynamics in this realm. Since ρ_S is diagonal, it follows (Supplement D) that $\sigma(t)$ is positive iff $\frac{d}{dt}|\rho_{ee}(t) - (\rho_0)_{ee}| \leq 0$, consistently with the interpretation of the relative entropy $\mathcal{S}(\rho_S||\rho_0)$ in (12b) as the entropic “distance” from equilibrium. Conversely, whenever the oscillatory $\rho_{ee}(t)$ drifts away from its initial or final equilibria, σ takes negative values (Fig. 1(b)).




FIG. 1: System and bath evolution as a function of time. Main panel: Excited-level population as a function of time for initially zero-temperature product state, followed by relaxation to quasi-equilibrium and then subjected to a series of measurements (vertical dashed lines). Measurements of finite duration ($\tau_k = 0.11/\omega_a$) (black line) results in somewhat larger heat-up than impulsive measurements (red line), but the dominant effect is the same for both. Observe the agreement between 2^{nd} order master equation, two-quanta exchange with a discrete bath (Suppl. A), and exact numerical solution for a discrete bath of 40 modes. (a) Relaxation rates, R_g , R_e as a function of time. (b) $\sigma(t)$ (negative of relative entropy rate of change). (c) Excitations as a function of t of the 40 modes in the two-quanta model. Parameters: $t_c = 10/\omega_a$, $\omega_0 = \omega_a$, $\gamma = 0.07\omega_a$.

In order to realistically model the repeated measurements, i.e. give them finite duration, we assume a smooth temporal profile of the coupling to the detector (Supplement A). The k -th measurement then occurs at time t_k and has a duration of τ_k . Figure 1(main panel) compares the population evolution via projective (impulsive) and finite-duration measurements with $\tau_k \sim 0.1/\omega_a$. Finite-duration measurements increase the Zeno-heating as compared to impulsive ones due to the extra energy supplied by the coupling to the detector. However, the




FIG. 2: Maximal system heating and cooling. Main panel: Maximum heating (lower half) and cooling (upper half) of the system, for different system- (x -axis) and bath- (y -axis) initial temperatures: $\alpha_S \neq \alpha_B$, where $\alpha_{S(B)} = 1/\beta_{S(B)}\hbar\omega_a$. (a) $R_g(t)$ and $R_e(t)$ (Eq. (6)) depicted as spectral overlaps of the relevant functions. (b) Example of a system experiencing first Zeno heating, then *oscillatory*-Zeno cooling. (c) Maximal Zeno heating (red) and subsequent maximal cooling (blue) as a function of common initial temperature of system and bath. Note the *critical temperature* for oscillatory-Zeno cooling. Parameters: $t_c = 10/\omega_a$, $\omega_0 = \omega_a/0.7$, $\gamma = 4.36\omega_a$. These effects can be *strongly magnified* by choosing other suitable parameters. (d) Possible experimental setup.

basic effect is seen (Fig. 1 – main panel) to be the same and is governed by $\langle H_{SB} \rangle$ change in Eq. (3). Counter-intuitively, finite-duration measurements are able to *increase the cooling*, despite the extra energy coupled in by the apparatus.

If we repeat this procedure often enough, the TLS will either increasingly heat up or cool down, upon choosing the time intervals Δt_k to coincide with either peaks or troughs of the ρ_{ee} oscillations, respectively. The minimal value of σ can also be progressively lowered with each measurement (Fig. 1(b)). Since consecutive measurements affect the bath and the system differently, they may acquire different temperatures, which then become the initial conditions for subsequent QZE heating or OZE cooling. The results are shown in Fig. 2 for both different (main panel) and common (Fig. 2(c)) temperatures of the system and the bath. Remarkably, the system may heat up solely due to the QZE, although the *bath is colder*, or cool down solely due to the OZE or AZE, although the *bath is hotter* (Fig. 2 – main panel). The bath may undergo changes in temperature and entropy too (Fig. 1(c)).

One experimental realization of these effects can involve atoms or molecules in a microwave cavity (Fig. 2(d)) with controllable finite-temperature coupling spectrum $G_T(\omega)$ centered at ω_0 . Measurements can be effected on such a TLS ensemble with resonance frequency ω_a in the microwave domain, at time intervals $\Delta t_k \sim 1/(\omega_0 \pm \omega_a)$, by an optical QND probe[8] at frequency $\omega_p \gg \omega_a, \omega_0$. The probe pulses undergo different Kerr-nonlinear phase shifts $\Delta\phi_e$ or $\Delta\phi_g$ depending on

the different symmetries (e.g., angular momenta) of $|e\rangle$ and $|g\rangle$. The relative abundance of $\Delta\phi_e$ and $\Delta\phi_g$ would then reflect the ratio $\rho_{ee}(t_k)/\rho_{gg}(t_k)$. Such QND probing may be performed with time-duration much shorter than ω_a^{-1} , i.e. $\omega_a\tau_k \ll 1$, without resolving the energies of $|e\rangle$ and $|g\rangle$.

Since non-selective measurements increase the Von-Neumann entropy of the detector ancillae, their entropic price precludes a “perpetuum mobile”, if closed-cycle operation is attempted. Yet, if our ancillae are laser pulses, they are only used once and we may progressively change the TLS ensemble thermodynamics by consecutive pulses, disregarding their entropic or energetic price. The practical advantage of the predicted anomalies is the possibility of *very rapid control* of cooling and entropy, which may be attained after several measurements at $t \geq \omega_a^{-1}$ and is only limited by the measurement rate. By contrast, conventional cooling requires much longer times, $t \gg t_c$, to reach thermal equilibrium.

The present findings establish a new link between frequent quantum measurements and nonequilibrium thermodynamical anomalies: heat and entropy rates of change with the “wrong” sign, as compared to their usual monotonic approach to equilibrium. These anomalies are determined by the oscillatory or *negative* values of the non-Markovian quantum relaxation rates at short times corresponding to large energy uncertainty. They reveal unfamiliar aspects of post-measurement quantum dynamics: AZE, which has been conceived as a means of *enhancing or accelerating* the initial-state change[12, 13, 14, 15], here can either *restore* the equilibrium state or further depart from it via cooling. These anomalies underscore the often forgotten fact that the system and the bath are *inseparable* (entangled)[6, 20], even under weak-coupling conditions, a fact that has profound implications on their short-time dynamics.

These results prompt further studies of a hitherto unexplored non-Markovian time domain where existing formulations of the second law of thermodynamics are inapplicable[2, 5], and neither is the common notion that heat always flows from hotter to colder ensembles. This domain may necessitate an in-depth scrutiny of quantum thermodynamical concepts. In particular, the need for temporal “coarse-graining” of entropy should be examined.

Acknowledgments

We acknowledge the support of ISF, GIF and EC (SCALA IP).

SUPPLEMENTARY INFORMATION

Supplement A. Measuring the Energy of a Two-Level System Interacting with a Bath: Dynamical Description

A.1 Hamiltonians and the measurement process

We consider the following Hamiltonians:

$$H(t) = H_{tot} + H_{SD}(t), \quad H_{tot} = H_S + H_B + H_{SB}. \quad (13)$$

Here H_{tot} pertains to the coupled system and bath and consists of:

$$H_S = \hbar\omega_a|e\rangle\langle e|, \quad (14)$$

$$H_B = \hbar \sum_{\lambda} \omega_{\lambda} a_{\lambda}^{\dagger} a_{\lambda}, \quad (15)$$

$$H_{SB} = \mathcal{S}\mathcal{B}, \mathcal{S} = |g\rangle\langle e| + |e\rangle\langle g|,$$

$$\mathcal{B} = \hbar \sum_{\lambda} \left(\kappa_{\lambda} a_{\lambda} + \kappa_{\lambda}^* a_{\lambda}^{\dagger} \right), \quad (16)$$

where $\mathcal{S}$ and $\mathcal{B}$ are the system and bath factors, respectively, in the system-bath interaction operator, $a_{\lambda}(a_{\lambda}^{\dagger})$ are the annihilation (creation) operators, and κ_{λ} is the matrix element of the weak coupling to bath mode λ .

The detector (ancilla) qubits have energy-degenerate states $|0\rangle_k, |1\rangle_k$ so that we may set the detector Hamiltonian to be zero

$$H_D = 0. \quad (17)$$

The time-dependent system-detector coupling (to the k th detector) has the form

$$H_{SD}(t) = \sum_k H_{SD,k} = \sum_{k=1}^M h_k(t) |e\rangle\langle e| (|0\rangle_{kk}\langle 0| + |1\rangle_{kk}\langle 1| - |0\rangle_{kk}\langle 1| - |1\rangle_{kk}\langle 0|). \quad (18)$$

where

$$h_k(t) = \frac{\pi}{4\tau_k} \left(\tanh^2 \left(\frac{t - t_k}{\tau_k} \right) - 1 \right) \quad (19)$$

is a smooth temporal profile of the system coupling to the detector qubits during the k -th measurement that occurs at time t_k and has a duration of τ_k .

This form of the single-measurement Hamiltonian $H_{SD,k}$ was chosen so that (dropping the index k and taking the measurement interval to be $[0, \tau]$ for simplicity):

$$e^{-i \int_0^{\tau} dt H_{SD}(t)/\hbar} = U_C. \quad (20)$$

where U_C denotes to the CNOT operation (with the k th detector qubit, the target qubit, denoted by the subscript D):

$$\begin{aligned} |g\rangle|0\rangle_D &\mapsto |g\rangle|0\rangle_D \\ |e\rangle|0\rangle_D &\mapsto |e\rangle|1\rangle_D \\ |g\rangle|1\rangle_D &\mapsto |g\rangle|1\rangle_D \\ |e\rangle|1\rangle_D &\mapsto |e\rangle|0\rangle_D. \end{aligned} \quad (21)$$

Since we take the initial state of the detector to be $|0\rangle$, only the first two rows play a role. If the measurement

duration τ is much shorter than the other time scales, then only H_{SD} is non-negligible, and the entire action of $H(t)$ during this time is well approximated by the CNOT operator U_C . This becomes exact in the impulsive limit $\tau \rightarrow 0$.

The measurement consists in letting the TLS interact with the detector (a degenerate TLS) via H_{SD} . The measurement outcomes are averaged over (for nonselective measurements), by tracing out the detector degree of freedom. The total effect on the system density-operator is:

$$\rho_S \mapsto \text{Tr}_D \{ U_C \rho_S \otimes |0\rangle_{DD}\langle 0| \} = |e\rangle\langle e| \rho_S |e\rangle\langle e| + |g\rangle\langle g| \rho_S |g\rangle\langle g| \quad (22)$$

i.e., the diagonal elements are unchanged, and the off-diagonals are erased. Since the TLS is entangled with the bath, the effect of the measurement in Eqs. (20) - (21) is:

$$\begin{aligned} \rho_{tot} &\mapsto \text{Tr}_D \{ U_C \rho_{tot} \otimes |0\rangle_{DD}\langle 0| \} \\ &= |e\rangle\langle e| \rho_{tot} |e\rangle\langle e| + |g\rangle\langle g| \rho_{tot} |g\rangle\langle g| \equiv \rho_{ee}^B |e\rangle\langle e| + \rho_{gg}^B |g\rangle\langle g|. \end{aligned} \quad (23)$$

A.2 Non-commutativity of H_{SD}, H_{SB} and the measurement-induced vanishing of $\langle H_{SB} \rangle$

Here we demonstrate the validity of Eq. (2) of the main text, based on the post-measurement increase in H_{tot} following the vanishing of $\langle H_{SB} \rangle$. We show that this effect *disappears* if $H_{SD}(t)$ commutes with H_{SB} and H_S (and hence with H_{tot}).

For the commutative case, we have:

$$\langle H_{tot}(\tau) \rangle = \text{Tr} \{ \rho_{SBD}(\tau) H_{tot} \} \quad (24)$$

$$= \text{Tr} \left\{ e^{-i \int_0^\tau dt [H_{tot} + H_{SD}(t)]} [\rho_{tot}(0) |0\rangle_D {}_D\langle 0|] \times e^{+i \int_0^\tau dt' [H_{tot} + H_{SD}(t')]} H_{tot} \right\} \quad (25)$$

$$= \text{Tr} \left\{ \rho_{tot}(0) |0\rangle_D {}_D\langle 0| e^{+i \int_0^\tau dt' [H_{tot} + H_{SD}(t')]} H_{tot} \times e^{-i \int_0^\tau dt [H_{tot} + H_{SD}(t)]} \right\} \quad (26)$$

$$= \text{Tr} \{ \rho_{tot}(0) H_{tot} \} \equiv \langle H_{tot}(0) \rangle, \quad (27)$$

where ρ_{SBD} is the state of the combined system, bath, and detector. The cyclic property of the trace was used in (26), and the commutativity of H_{SD} and H_{tot} in (27).

Compare this now to our *non-commutative* model (Eqs. (16) - (20))

$$e^{-i \int_0^\tau dt H_{SD}(t)} |0\rangle_D = U_C |0\rangle_D = |1\rangle_D |e\rangle \langle e| + |0\rangle_D |g\rangle \langle g|. \quad (28)$$

Since H_{SD} in Eq. (18) commutes with H_S , we may consider the evolution of $\langle H_{SB}(\tau) \rangle$, rather than $\langle H_{tot}(\tau) \rangle$. In the impulsive limit ($\tau \rightarrow 0$), we can drop H_{tot} in the exponent of Eq. (26), and then use the LHS of (28) to obtain:

$$\langle H_{SB}(\tau) \rangle = \text{Tr} \left\{ \rho_{tot}(0) {}_D\langle 0| U_C^\dagger H_{SB}(0) U_C |0\rangle_D \right\}. \quad (29)$$

Finally, using the RHS of (28) and (16), we get:

$${}_D\langle 0| U_C^\dagger H_{SB}(0) U_C |0\rangle_D = 0 \rightarrow \langle H_{SB}(\tau) \rangle = 0. \quad (30)$$

This expresses the vanishing of $\text{Tr} \{ \rho_{tot}(\tau) H_{SB} \}$ due to the diagonality of $\rho_{tot}(\tau)$ with respect to S (Suppl. B). Since $H_D = 0$, the detector mean energy is not affected by the CNOT action.

Hence, the measurement-induced interruption of the mean interaction energy, $\langle H_{SB}(\tau) \rangle = 0$, and the resulting $\langle H_S \rangle + \langle H_B \rangle$ changes in Eq.(2) of the main text have a quantum mechanical origin: the non-commutativity of H_{SB} and H_{SD} .

A.3 Two-quanta approximation

Here we assume that the system and bath, governed by Eqs. (13) - (18) above, were in their respective ground states prior to their interaction onset at $t = 0$, followed by a measurement at time t_k . As shown in Suppl. B, one should allow for arbitrary excitations of the system and bath leading to an infinite hierarchy of coupled equations for the populations of $|e\rangle, |g\rangle$ and mode excitation numbers. Here, for simplicity, we curtail this hierarchy, as is justified at short times. The wave-function driven by $H(t)$ (Eq. (13)) acquires the following form, by allowing the system+bath to receive or give away only 0 or 2 excitations (the lowest two orders of the hierarchy expansion) through the coupling to the detector:

$$|\psi(t)\rangle = \sum_{l=0}^{2^M-1} |\psi^{(l)}(t)\rangle |b_l\rangle \quad (31)$$

$$|\psi^{(l)}(t)\rangle = \tilde{\alpha}_{g,0}^{(l)}(t) |g\rangle \bigotimes_{\lambda} |0\rangle_{\lambda} + \sum_{\lambda} \tilde{\alpha}_{e,\lambda}^{(l)}(t) |e\rangle |1\rangle_{\lambda} \bigotimes_{\lambda' \neq \lambda} |0\rangle_{\lambda'} \quad (32)$$

$$+ \sum_{\lambda} \tilde{\alpha}_{g,\lambda}^{(l)}(t) |g\rangle |2\rangle_{\lambda} \bigotimes_{\lambda' \neq \lambda} |0\rangle_{\lambda'} + \sum_{\lambda < \lambda'} \tilde{\alpha}_{g,\lambda,\lambda'}^{(l)}(t) |g\rangle |1\rangle_{\lambda} |1\rangle_{\lambda'} \bigotimes_{\lambda'' \neq \lambda, \lambda'} |0\rangle_{\lambda''} \quad (33)$$

where b_l is the binary representation of l , labelling the detector qubits. We transform to the frame where am-

plitudes are defined by:

$$\tilde{\alpha}_{g,0}^{(l)} = \alpha_{g,0}^{(l)} \quad (34)$$

$$\tilde{\alpha}_{e,\lambda}^{(l)} = e^{-i\omega_a t - i\omega_{\lambda} t} \alpha_{e,\lambda}^{(l)} \quad (35)$$

$$\tilde{\alpha}_{g,\lambda}^{(l)} = e^{-i2\omega_{\lambda} t} \alpha_{g,\lambda}^{(l)} \quad (36)$$

$$\tilde{\alpha}_{g,\lambda,\lambda'}^{(l)} = e^{-i(\omega_{\lambda} + \omega_{\lambda'}) t} \alpha_{g,\lambda,\lambda'}^{(l)}. \quad (37)$$

Using the Schrödinger equation, and integrating explicitly for $\alpha_{g,0}^{(l)}$, $\alpha_{g,\lambda}^{(l)}$ and $\alpha_{g,\lambda,\lambda'}^{(l)}$, we obtain the following integro-differential matrix equation:

$$\dot{\alpha}_e^{(l)}(t) = -R(t)\alpha_e^{(l)}(t) - i\mathbf{f}^{(l)}(t) - i \sum_{k=1}^M h_k(t) \left(\alpha_e^{(l)} - \alpha_e^{(Q_k(l))} \right) \quad (38)$$

$$R_{\lambda,\lambda'}(t) = \int_0^t dt' \left\{ \kappa_{\lambda}^* \kappa_{\lambda'} \left[e^{-i(\omega_{\lambda'} t' - \omega_{\lambda} t)} + e^{-i(\omega_{\lambda'} t - \omega_{\lambda} t')} \right] + \delta_{\lambda,\lambda'} \sum_{\lambda''} \kappa_{\lambda''}^* \kappa_{\lambda'} e^{-i\omega_{\lambda''}(t-t')} \right\} e^{-i\omega_{\lambda}(t'-t)} \quad (39)$$

$$f_{\lambda}^{(l)}(t) = \kappa_{\lambda}^* e^{i\omega_{\lambda} t + i\omega_{\lambda} t} \alpha_{g,0}^{(l)}(0) + \sqrt{2} \kappa_{\lambda} e^{i\omega_{\lambda} t - i\omega_{\lambda} t} \alpha_{g,\lambda}^{(l)}(0) + \sum_{\lambda \neq \lambda'} \kappa_{\lambda'} e^{i\omega_{\lambda} t - i\omega_{\lambda'} t} \alpha_{g,\lambda,\lambda'}^{(l)}(0) \quad (40)$$

where $\alpha_e^{(l)}(t) = \{\alpha_{e,\lambda}^{(l)}\}^T$, and $Q_k(l)$ is the decimal representation of $\{b_1, \dots, 1-b_k, \dots, b_l\}$, describing the flipping of the k th detector qubit. This flipping occurs within the $H_{SD}(t)$ activation interval defined in Eq. (38) by $h_k(t)$ (Eq.(19)).

Performing a brief measurement of the system at time t_k (according to Eqs. (38)-(40)) “splits” the subsequent evolution into two paths: (i) detection of the excited state with probability $P^{(e)}(t_k) = \sum_{\lambda} |\alpha_{e,\lambda}(t_k)|^2$, and $\alpha_{g,0}^{(e)}(t_k + \epsilon) = \alpha_{g,\lambda}^{(e)}(t_k + \epsilon) = \alpha_{g,\lambda,\lambda'}^{(e)}(t_k + \epsilon) = 0$, where the superscript (e) denotes the excited-state outcome; (ii) detection of the ground state with probability $P^{(g)}(t_k) = 1 - P^{(e)}(t_k)$ and $\alpha_{e,\lambda}(t_k + \epsilon)^{(g)} = 0$, the superscript (g) denoting the ground-state. These subsequent independent evolutions destroy the system-bath correlations, and can give rise to the phenomena of Zeno heating and OZE cooling described in the text.

The plots in Fig. 1 of the text confirm the adequacy of the present two-quanta approximation in describing

the approach to equilibrium and the measurement effects, compared to exact numerical simulations (Suppl. C) or the second-order master-equation approach.

Supplement B. Bath-System Entanglement Near Thermal Equilibrium: Pre- and Post-Measurement States

The eigenstates of the total Hamiltonian H_{tot} (Eq. (13)) exhibit entanglement between the system and bath, due to the interaction term H_{SB} . The same is true of thermal states ($Z^{-1}e^{-\beta H_{tot}}$), at least at low temperatures and for weak coupling.

Likewise, if the system and bath are initially in a factorizable eigenstate of $H_0 = H_S + H_B$ and are subsequently exposed to the total Hamiltonian, including the interaction term, they will evolve into entangled system-bath states, at any temperature. If the interaction is turned on adiabatically, an H_0 eigenstate may be expected to evolve asymptotically into an eigenstate of H_{tot} .

The aim of this Supplement is to prove the assertion that in all these cases, the following properties of the joint state of the system and bath obtain:

- (i) ρ_S is always diagonal (in the H_S basis), which implies in turn $Tr \{\rho_{tot} H_{SB}\} = 0$.
- (ii) The first moment of the bath excitation or deexcitation operator vanishes, causing the initial vanishing of the time derivative of ρ_S immediately after a measurement (the Zeno effect).
- (iii) There is short-time post-measurement (Zeno) heating.

B.1 Pre-measurement evolution towards equilibrium

We shall work in the interaction picture:

$$V_I(t) \equiv e^{+iH_0(t-t_0)} H_{SB} e^{-iH_0(t-t_0)} = (e^{-i\omega_a(t-t_0)} |e\rangle\langle g| + e^{+i\omega_a(t-t_0)} |g\rangle\langle e|) \sum_k (e^{i\omega_0(t-t_0)} \kappa_k^* a_k^\dagger + \kappa_k a_k e^{-i\omega_0(t-t_0)}), \quad (41)$$

$$|\Psi_I(t)\rangle \equiv e^{iH_0(t-t_0)} |\Psi(t)\rangle = U(t, t_0) |\Psi_I(t_0)\rangle, \quad (42)$$

$$U(t, t_0) = 1 + \sum_{n=1}^{\infty} (-i)^n \int_{t_0}^t dt_1 \int_{t_0}^{t_1} dt_2 \cdots \int_{t_0}^{t_{n-1}} dt_n V_I(t_1) V_I(t_2) \cdots V_I(t_n) \equiv \sum_{n=0}^{\infty} O_n(t). \quad (43)$$

Let us denote the joint eigenstates of H_B and $\hat{N}$ (the total number operator) by:

$$|\mathbf{n}\rangle \equiv \mathcal{N}_{\mathbf{n}} \prod_j (a_{k_j}^\dagger)^{n_j} |0\rangle, \quad (44)$$

where $\mathbf{n} \equiv \{n_j\}_j$ and $\mathcal{N}_{\mathbf{n}}$ is the appropriate normalization constant.

Consider first $|\Psi(t_0)\rangle = |\mathbf{n}\rangle \otimes |g\rangle$, where $|\mathbf{n}\rangle$ has the $\hat{N}$ eigenvalue n_{tot} . We note that $V_I(t)$ (for any t) has the effect of flipping the H_S state and transforming the bath state into a sum of states with one more or one less excitation. Therefore the even terms ($O_{2m}(t)|\Psi(t_0)\rangle$) in the perturbation expansion of $|\Psi_I(t)\rangle$, Eq. (43) (counting the 1 as the zeroth term!), are superpositions of states with excitation numbers $n_{tot} + \text{even}$, multiplied by $|g\rangle$, while the odd ones ($O_{2m+1}(t)|\Psi(t_0)\rangle$) are superpositions of states with $n_{tot} + \text{odd}$, multiplied by $|e\rangle$. Let us denote the sum of the even terms of the series by $|B^{\text{even}}\rangle|e\rangle$ and that of the odd terms as B^{odd} , then:

$$\begin{aligned} |\Psi_I(t)\rangle &= U(t, t_0)|\mathbf{n}, g\rangle = |B_{\mathbf{n},g}^{\text{even}}(t)\rangle \otimes |g\rangle + |B_{\mathbf{n},g}^{\text{odd}}(t)\rangle \otimes |e\rangle \\ &\equiv |\Psi_{\mathbf{n},g}(t)\rangle. \end{aligned} \quad (45)$$

Here B^{even} (respectively, B^{odd}) is a sum of $\hat{N}$ -eigenstates with eigenvalues differing from $\hat{N}$ by even (respectively, odd) numbers.

If the initial state is of the form $|\Psi(t_0)\rangle = |\mathbf{n}\rangle \otimes |e\rangle$, the time-evolved state is:

$$\begin{aligned} |\Psi_I(t)\rangle &= U(t, t_0)|\mathbf{n}, e\rangle = |B_{\mathbf{n},e}^{\text{even}}(t)\rangle \otimes |e\rangle + |B_{\mathbf{n},e}^{\text{odd}}(t)\rangle \otimes |g\rangle \\ &\equiv |\Psi_{\mathbf{n},e}(t)\rangle. \end{aligned} \quad (46)$$

ndequation

Now consider the initial condition that the system and bath are in a Gibbs state of H_0 , with any inverse temperature β :

$$\begin{aligned} \rho_{tot}(t_0) &= Z_{tot}^{-1} e^{-\beta H_0} = Z_S^{-1} e^{-\beta H_S} Z_B^{-1} e^{-\beta H_B} \\ &= Z_{tot}^{-1} \sum_{\mathbf{n}} \left\{ e^{-\beta(\omega_g + \omega_{\mathbf{n}})} |\mathbf{n}, g\rangle \langle \mathbf{n}, g| + e^{-\beta(\omega_e + \omega_{\mathbf{n}})} |\mathbf{n}, e\rangle \langle \mathbf{n}, e| \right\} \end{aligned} \quad (47)$$

This initial state evolves at time $t \gg t_0$ into:

$$\begin{aligned} \rho_{tot}(t) &= Z_{tot}^{-1} \sum_{\mathbf{n}} \left\{ e^{-\beta(\omega_g + \omega_{\mathbf{n}})} |\Psi_{\mathbf{n},g}(t)\rangle \langle \Psi_{\mathbf{n},g}(t)| \right. \\ &\quad \left. + e^{-\beta(\omega_e + \omega_{\mathbf{n}})} |\Psi_{\mathbf{n},e}(t)\rangle \langle \Psi_{\mathbf{n},e}(t)| \right\} \end{aligned} \quad (48)$$

The resulting ρ_{tot} has off-diagonal $|e\rangle\langle g|$ and $|g\rangle\langle e|$ elements by virtue of Eqs. (45) and (46).

Assuming that the adiabatic theorem can be applied (despite the *initial* degeneracy of the eigenstates), we have under adiabatic switching on of H_{SB} :

$$\rho_{tot}(t \rightarrow \infty) = \sum_{\mathbf{n}} \sum_{m=g,e} e^{-\beta E_{\mathbf{n},m}} |\Psi_{\mathbf{n},m}(t)\rangle \langle \Psi_{\mathbf{n},m}(t)|, \quad (49)$$

$E_{\mathbf{n},m}$ being the eigenvalues of H_0 .

This state has Gibbsian form, but with the original Boltzmann weights. However, these weights are irrelevant for the purity of the state.

B.2 Diagonality of ρ_S and post-measurement vanishing of its first order derivative

We wish to establish the diagonality of ρ_S before ($t \leq 0$) and after ($t \geq \tau$) the measurement, and the vanishing of $\dot{\rho}_S$ immediately after the measurement ($t = \tau$). Due to the post-measurement vanishing of the off-diagonal elements of ρ_{tot} (Eq. (24), its derivative immediately after the measurement, $\dot{\rho}_S(\tau)$, has the form:

$$\dot{\rho}_S(\tau) = -i \left(e^{-i\omega_a\tau} |e\rangle\langle g| - e^{+i\omega_a\tau} |g\rangle\langle e| \right) \text{Tr}_B \left\{ \mathcal{B} (\rho_{gg}^B - \rho_{ee}^B) \right\}. \quad (50)$$

For $\rho_{tot}(t) = |\Psi_{\mathbf{n},g}(t)\rangle \langle \Psi_{\mathbf{n},g}(t)|$ (Eq. (45)), we then have

$$\begin{aligned} (\rho_S)_{eg}(t) &= \langle e | \rho_S(t) | g \rangle = \text{Tr}_B \langle e | \rho_{tot}(t) | g \rangle \\ &= \langle B_{\mathbf{n},e}^{\text{even}}(t) | B_{\mathbf{n},e}^{\text{odd}}(t) \rangle = 0 \end{aligned} \quad (51)$$

Hence, ρ_S is diagonal at any time t .

In addition, we have, by virtue of (50) at time $t = \tau$

$$\begin{aligned} \dot{\rho}_S(\tau) &\propto \text{Tr}_B \left\{ \mathcal{B} \langle e(g) | \rho_{tot} | e(g) \rangle \right\} \\ &= \langle B_{\mathbf{n},e}^{\text{even(odd)}}(\tau) | \mathcal{B} | B_{\mathbf{n},e}^{\text{even(odd)}}(\tau) \rangle = 0 \end{aligned} \quad (52)$$

The same argument goes through upon permuting $e \leftrightarrow g$ everywhere for $\rho_{tot} = |\Psi_{\mathbf{n},e}(t)\rangle \langle \Psi_{\mathbf{n},e}(t)|$ (Eq. (46)).

By linearity, using (51) and (52), the diagonality of $\rho_S(t)$ and the vanishing of $\dot{\rho}_S$ immediately after the measurement are satisfied for (48) and (49).

B.3 Second derivative positivity at arbitrary time

For the *factorizable* thermal state,

$$\rho_{tot} = Z^{-1} e^{-\beta H_0} = Z_B^{-1} e^{-\beta H_B} Z_S^{-1} e^{-\beta H_S}, \quad (53)$$

we have:

$$\begin{aligned} \rho_{ee}^B &\equiv \langle e | \rho_{tot} | e \rangle = \langle e | Z_S^{-1} e^{-\beta H_S} | e \rangle Z_B^{-1} e^{-\beta H_B} = (\rho_S)_{ee} \rho_B \\ &\text{(and } e \leftrightarrow g \text{)}. \end{aligned} \quad (54)$$

For this ρ_{tot} , the second derivative of ρ_S immediately after the measurement is (cf. Eq. (24))

$$\ddot{\rho}_S(\tau) = 2\sigma_z \text{Tr}_B \left\{ \hat{\mathcal{B}}^2 (\rho_{gg}^B - \rho_{ee}^B) \right\}. \quad (55)$$

The scalar factor is positive:

$$\text{Tr}_B \left\{ \hat{\mathcal{B}}^2 (\rho_{gg}^B - \rho_{ee}^B) \right\} = \text{Tr}_B \left\{ \hat{\mathcal{B}}^2 \rho_B \right\} \left((\rho_S)_{gg} - (\rho_S)_{ee} \right) > 0, \quad (56)$$

where we have used $\text{Tr}_B \{ \rho_{gg}^B \} = (\rho_S)_{gg}$ which follows from the definition (Eq. (24)): $\rho_{ee}^B = \langle e(g) | \rho_{tot} | e(g) \rangle$. The first factor in (56) is positive by virtue of the positivity of the operator $\hat{\mathcal{B}}^2$ ($\hat{\mathcal{B}}$ being Hermitian), and the second is positive iff there is no population inversion for the TLS.

The combined (system- and bath-) equilibrium state satisfies:

$$\rho_{tot} = Z^{-1} e^{-\beta H_{tot}} = \rho_{tot} = Z^{-1} e^{-\beta(H_0 + O(H_{SB}^2))}. \quad (57)$$

Thus, for sufficiently weak coupling, Eq. (54) dominates.

We have made no use of the equality of the system and bath temperatures. The argument goes through unchanged for $\rho_{tot} = Z_B^{-1} e^{-\beta_B H_B} Z_S^{-1} e^{-\beta_S H_S}$.

C. Exact Numerical Simulations

The numerical calculations have been done by the Multi-Configuration Time-Dependent Hartree (MCTDH) approach[19, 21]. With this approach, very large system-bath wave functions (with more than 100 degrees of freedom) can be propagated in time with very high precision. In order to treat finite temperatures we have sampled the Boltzmann operator in an efficient way, by the random-phase thermal wave function technique [6].

D. Entropy Dynamics

We may always *define* the entropy of ρ_S *relative* to its equilibrium state ρ_0 and the negative of its rate of change:

$$\mathcal{S}(\rho_S(t)||\rho_0) \equiv \text{Tr}\{\rho_S(t) \ln \rho_S(t)\} - \text{Tr}\{\rho_S(t) \ln \rho_0\} \quad (58)$$

$$\sigma(t) \equiv -\frac{d}{dt} \mathcal{S}(\rho_S(t)||\rho_0). \quad (59)$$

Only in the Markovian realm, σ thus defined is identified as the ‘entropy production rate’, i.e., the net rate of change of the entropy of the system, after deducting the entropy change due to the exchange of heat $\mathbf{Q}$ with the bath at temperature $\mathbf{T}$ (in the absence of external work) [3]:

$$\sigma(t) = \frac{d\mathcal{S}(\rho_S)}{dt} - \frac{1}{\mathbf{T}} \frac{d\mathbf{Q}}{dt}. \quad (60)$$

where $\mathcal{S}(\rho_S)$ is the *absolute* Von-Neumann entropy of the system.

It was proven by Lindblad [5] that under any proper (completely positive) quantum dynamical map, $\mathbf{M}$, the *relative* entropy, Eq.(58) cannot increase: $\mathcal{S}(\mathbf{M}\rho_S||\mathbf{M}\rho_0) \leq \mathcal{S}(\rho_S||\rho_0)$. It then follows that under Markovian evolution, which is described by such a map,

$$\sigma(t) \geq 0. \quad (61)$$

This is a statement of the second law of thermodynamics[2].

The quantity $\sigma(t)$ defined in Eq. (59) (which only in the Markovian case is identified as the entropy production rate of the TLS) has the following form, with $(\rho_0)_{ee}$ as the long-time (equilibrium) excitation probability

$$\sigma = -\dot{\rho}_{ee} \ln \left[\frac{\rho_{ee}(1 - (\rho_0)_{ee})}{(\rho_0)_{ee}(1 - \rho_{ee})} \right]. \quad (62)$$

Thus, $\sigma(t)$ is positive iff $\frac{d}{dt} |\rho_{ee}(t) - (\rho_0)_{ee}| \leq 0$, consistently with the interpretation of the relative entropy $\mathcal{S}(\rho_S||\rho_0)$ in (58) as a kind of “distance” from equilibrium. Conversely, whenever the oscillatory $\rho_{ee}(t)$ drifts away from its initial or final equilibria, σ must take negative values (Fig. 1(b)).

The condition for $\sigma < 0$, amounts to:

$$R_e(t)\rho_{ee} - R_g(t)\rho_{gg} < 0 \quad \text{and} \quad \ln \frac{\rho_{ee}}{\rho_{gg}} \quad (63)$$

Lindblad’s theorem on the effect of quantum maps on the relative entropy[5] seems at first sight to contradict our results. The resolution of this apparent paradox is that the non-Markov evolution of $\rho_S(t)$ is not described by a map at all! This can be seen (Figs. 1,2b) from the fact that the density matrix oscillates and goes through the *same value* more than once, with differing subsequent evolution. Hence, $\mathbf{M}(0,t) : \rho_S(0) \mapsto \rho_S(t)$ is a well defined completely positive map, but it is not invertible: different values of $\rho_S(0)$ may evolve into the same $\rho_S(t)$ for some particular time t . This bears likeness to the findings in Ref. [20] in a different context.

-
- [1] LD Landau and EM Lifshitz. *Statistical Physics 3rd edn, part 1*. Pergamon Press, 1980.
 - [2] H. Spohn. Entropy production for quantum dynamical semigroups. *J. Math. Phys.*, 19:1227, 1978.
 - [3] R. Alicki. The quantum open system as a model of the heat engine. *J. Phys. A*, 12:L103, 1979.
 - [4] C. Jarzynski. Nonequilibrium equality for free energy differences. *Phys. Rev. Lett.*, 78:2690, 1997.
 - [5] G. Lindblad. Expectations and entropy inequalities for finite quantum systems. *Comm. Math. Phys.*, 39(2):111–

119, 1974.

- [6] D. Gelman and R. Kosloff. Simulating dissipative phenomena with a random phase thermal wavefunctions, high temperature application of the Surrogate Hamiltonian approach. *Chem. Phys. Lett.*, 381(1):129–138, 2003.
- [7] Wojciech Hubert Zurek. Decoherence, einselection, and the quantum origins of the classical. *Rev. Mod. Phys.*, 75(3):715–775, May 2003.
- [8] V.B. Braginsky and F.Y. Khalili. *Quantum Measurement*. Cambridge University Press, 1995.

- [9] SM Barnett. Turning on quantum taps. *Nature(London)*, 362(6416):113–113, 1993.
- [10] S. Haroche and J.M. Raimond. *Exploring the quantum: atoms, cavities, and photons*. Oxford University Press, 2006.
- [11] B. Misra and E. C. G. Sudarshan. Zeno’s paradox in quantum theory. *J. Math. Phys.*, 18:756–763, 1977.
- [12] A. M. Lane. Decay at early times - larger or smaller than the golden rule. *Phys. Lett.*, 99A:359–360, 1983.
- [13] A. G. Kofman and G. Kurizki. Acceleration of quantum decay processes by frequent observations. *Nature (London)*, 405:546–550, 2000.
- [14] P. Facchi and S. Pascazio. Quantum Zeno and inverse quantum Zeno effects. *Progress in Optics*, 42:147, 2001.
- [15] A. G. Kofman and G. Kurizki. Unified theory of dynamically suppressed qubit decoherence in thermal baths. *Phys. Rev. Lett.*, 93:130406, 2004.
- [16] L. S. Schulman and B. Gaveau. Ratcheting up energy by means of measurement. *Phys. Rev. Lett.*, 97:240405, 2006.
- [17] Marlan O. Scully. Extracting work from a single thermal bath via quantum negentropy. *Phys. Rev. Lett.*, 87(22):220601, Nov 2001.
- [18] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg. *Atom-Photon Interactions*. Wiley, New York, 1992.
- [19] M. Nest and H.D. Meyer. Dissipative quantum dynamics of anharmonic oscillators with the multiconfiguration time-dependent Hartree method. *J. Chem. Phys.*, 119:24, 2003.
- [20] Peter Stelmachovic and Vladimir Buzek. Dynamics of open quantum systems initially entangled with environment: Beyond the kraus representation. *Phys. Rev. A*, 64(6):062106, Nov 2001.
- [21] MH Beck, A Jäckle, GA Worth, and H.-D. Meyer. *Phys. Rep.*, 324(1):1, 2000.